

Technical Data Sheet

Human CD32b/c /Fc gamma RIIB (C-His-Avi)

Catalog Number: 820501, 820502

Size: 25 ug, 100 ug

Target Name: CD32b/c, FCGR2B, C, FcRII-b, c, CD32, FCG2, IGFR2

Regulatory Status: RUO

Product Details

Application: ELISA, BLI

Format: Liquid, Purified

Expression Host: CHO

Species: Human

Sources: Recombinant CD32b/c /Fc gamma RIIB (Ala46-Pro217) with C-terminus Avi-His-tag is expressed in CHO cell.

Accession Number: P31994

Molecular Weight: The protein has a predicted molecular weight of 23.5 kDa. Under DTT-reducing conditions, it migrates at approximately 35 kDa on SDS-PAGE.

Affinity Tag: C-Avi-His

Purity: >95% based on SDS-PAGE under reducing condition

Formulation: 1xPBS buffer, pH7.4, 0.22 µm filtered

Endotoxin level: Not tested

Protein Concentration: 25µg size is bottled at 0.2mg/mL concentration. 100 µg size is supplied at a lot-specific concentration.


Storage and Handling: Briefly centrifuge the vial upon receipt. An unopened vial can be stored at 4°C for up to 2 weeks, or at -20°C or below for up to six months. The protein may be further diluted to 0.1 mg/mL using 0.22 µm-filtered PBS buffer (pH 7.4). For long-term storage, the diluted stock solution should be aliquoted and stored at ≤ -70°C to minimize freeze-thaw cycles. If additional dilution is required, carrier proteins such as FBS or BSA should be added to maintain protein stability.

Background Information

CD32B (FCGR2B) and CD32C (FCGR2C) are 40 kDa, type I transmembrane proteins belonging to the Ig superfamily of low-affinity IgG Fc receptors. CD32B contains an immunoreceptor tyrosine-based inhibition motif (ITIM) in its cytoplasmic tail, while CD32C has an immunoreceptor tyrosine-based activation motif (ITAM). Both receptors can bind monomeric IgG and IgG complexes, and are expressed on B cells, monocytes, macrophages, granulocytes, platelets, and mast cells. CD32B primarily acts as a negative regulator, inhibiting immune cell activation, proliferation, endocytosis, phagocytosis, and degranulation. In contrast, CD32C has an activating function, contributing to immune responses. Together, they play crucial roles in regulating immune cell functions and maintaining immune homeostasis. CD32B and CD32C share almost identical extracellular domain. The aa sequence of Fc gamma RIIB and Fc gamma RIIC are the same for the

region expressed (Ala46-Pro217)

Product Data



Human CD32b/c /Fc gamma RIIB (C-His-Avi) on SDS-PAGE under reducing condition (P+) and non-reducing condition (P-). The purity of this protein appears to be greater than 95%.