

Human GM2A Protein (C-His)

Catalog Number:	601401, 601402
Size:	25 ug, 100 ug
Target Name:	GM2A, Ganglioside GM2 activator
Regulatory Status:	RUO


PRODUCT DETAILS

Application:	ELISA, BLI
Format:	Liquid, Purified
Expression Host:	HEK293
Species:	Human
Accession Number:	P17900
Sources:	Recombinant human GM2A (His24-Ile193) with C-terminus His tag was expressed in 293 Cells
Molecular Weight:	This protein has a predicted molecular weight of 20 kDa. Under DTT-reducing conditions, the protein migrates at approximately 24 kDa on SDS-PAGE.
Affinity Tag:	C-His
Purity:	>95% based on SDS-PAGE under reducing condition
Formulation:	1xPBS with 300mM NaCl, pH7.4, 0.22 µm filtered
Endotoxin level:	Not tested
Protein Concentration:	25µg size is bottled at 0.2mg/mL concentration. 100 µg size is supplied at a lot-specific concentration.
Storage and Handling:	Briefly centrifuge the vial upon receipt. An unopened vial can be stored at 4°C for up to 2 weeks, or at -20°C or below for up to six months. The protein may be further diluted to 0.1 mg/mL using 0.22 µm-filtered PBS buffer (pH 7.4). For long-term storage, the diluted stock solution should be aliquoted and stored at ≤ -70°C to minimize freeze-thaw cycles. If additional dilution is required, carrier proteins such as FBS or BSA should be added to maintain protein stability.

BACKGROUND INFORMATION

GM2 ganglioside activator (GM2A) is a lipid transfer protein of the ML domain family that facilitates the degradation of ganglioside GM2 by acting as a substrate-specific cofactor for β -hexosaminidase A. GM2A extracts GM2 from membranes and presents it in soluble form to β -hexosaminidase A for cleavage, enabling conversion to GM3. It can bind several single-chain phospholipids and fatty acids and shows some calcium-independent phospholipase activity. Mutations in GM2A cause GM2-gangliosidosis type AB, a Tay-Sachs disease variant characterized by impaired GM2 degradation and lipid accumulation.

PRODUCT DATA



Purified Human GM2A (His24-Ile193, with C-terminus His tag, 293 expressed) final product on SDS-PAGE under reducing (P+) conditions. The purity of Human GM2A appears to be greater than 95%.

This product is supplied subject to the terms and conditions at www.innocyto.com/web/terms.php and may only be used as provided in the stated terms. Products are for Research Use Only.