

Anti-Mouse CD49b (pan-NK, integrin α 2) Antibody

Catalog Number:	202701, 202702
Size:	100 ug, 25 tests
Target Name:	CD49b, Integrin alpha 2 chain, ITGA2
Regulatory Status:	RUO

PRODUCT DETAILS

Clone:	DX5_R
Application:	Flow Cytometry
Reactivity:	Mouse
Format:	Purified
Isotype:	Mouse IgG2a IgM like
Antibody Type:	Monoclonal
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide
Protein Concentration:	0.5 mg/mL
Storage&Handling:	The antibody solution should be stored between 2°C and 8°C
Recommended Usage:	For flow cytometric staining, it is recommended to use less than 0.2 μ g of this reagent per 0.5-1.0 million cells in a 100 μ L volume. Optimal reagent performance should be determined by titration for each specific application.
Isotype Control:	301501

BACKGROUND INFORMATION

Mouse CD49b, also known as integrin α 2 (Itga2), is a cell surface adhesion receptor best known in immunology as a defining marker of natural killer (NK) cells in mice. CD49b pairs with the integrin β 1 subunit (CD29) to form the α 2 β 1 integrin, also referred to as very late antigen-2 (VLA-2). Through this heterodimeric complex, CD49b mediates cell adhesion, migration, and signaling, supporting immune cell interactions with the extracellular matrix and other cells.

Structurally, CD49b is a type I transmembrane glycoprotein with a large extracellular domain, a single transmembrane helix, and a short cytoplasmic tail. The extracellular region contains an inserted (I) domain, also called an A domain, which is responsible for ligand binding and requires divalent cations such as Mg²⁺ or Mn²⁺ for activity. Like other integrins, CD49b undergoes conformational changes that regulate ligand affinity and enable bidirectional "inside-out" and "outside-in" signaling. The cytoplasmic tail lacks intrinsic enzymatic activity but associates with adaptor proteins that link the receptor to the actin cytoskeleton.

The primary ligands for mouse CD49b are extracellular matrix proteins, most notably collagen types I, II, and IV, as well as laminin. Binding to these ligands enables NK cells and other CD49b-expressing cells to adhere to tissue matrices and migrate within peripheral tissues. Through α 2 β 1 integrin signaling, CD49b contributes to cellular activation, survival, and cytotoxic function,


particularly in tissue-resident or tissue-infiltrating immune populations.

CD49b has important implications in disease and immune regulation. In mouse models, CD49b expression is widely used to identify and study NK cells and NKT cell subsets, including in cancer, infection, and autoimmune disease research. Altered $\alpha2\beta1$ integrin signaling has been linked to dysregulated immune cell trafficking, chronic inflammation, and fibrotic processes. In cancer models, CD49b-positive NK cells play a key role in antitumor immunity through their ability to localize to tumor tissues and mediate cytotoxic responses.


Therapeutically, mouse CD49b is primarily leveraged as a biomarker and experimental targeting handle rather than a direct clinical target. Antibodies against CD49b (such as DX5) are widely used to identify, isolate, or deplete NK cells in preclinical studies. Insights gained from CD49b-defined NK cell biology inform the development of NK cell-based immunotherapies and strategies aimed at enhancing immune cell trafficking and function, underscoring the translational relevance of this integrin in immunology research.

The HM α 2 antibody has been shown to be useful for partially blocking CD49b mediated interactions with collagen. Additionally, this antibody blocks staining of splenic NK cells by the monoclonal antibody DX5.

PRODUCT DATA



Mouse splenocytes were stained with FITC Anti-mouse NK1.1 clone PK136 and either purified Anti-Mouse CD49b clone DX5-R (right panel) or an isotype control (left panel).



Mouse splenocytes were stained with purified Anti-Mouse CD49b clone DX5-R (color-filled histogram) or an isotype control (gray histogram), followed by APC conjugated secondary antibody.

This product is supplied subject to the terms and conditions at www.innocyto.com/web/terms.php and may only be used as provided in the stated terms. Products are for Research Use Only.