

## Technical Data Sheet

### APC Anti-Human CD27

**Catalog Number:** 106707, 106708

**Size:** 25 tests, 100 tests

**Target Name:** CD27, TNFRSF7, S152, T14

**Regulatory Status:** RUO

---

#### Product Details

**Clone:** O323

**Application:** FC

**Reactivity:** Human

**Format:** APC

**Isotype:** Mouse IgG1

**Antibody Type:** Monoclonal

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA

**Protein Concentration:** Supplied at a lot-specific concentration.

**Storage&Handling:** The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.

**Recommended Usage:** For flow cytometric staining, it is recommended to use 5 uL of this reagent per 0.5-1.0 million cells in a 100  $\mu$ L volume. Optimal reagent performance should be determined by titration for each specific application.

**Excitation Laser:** Red Laser (633 nm)


---

#### Background Information

CD27 is a 50–55 kDa type I transmembrane glycoprotein, also known as S152 and T14, and is a lymphocyte-specific member of the tumor necrosis factor receptor (TNFR) superfamily, which also includes CD30 and CD40. Members of this superfamily are broadly involved in the regulation of cell proliferation, differentiation, and apoptosis. Unlike many other TNFR/TNF family members, expression of CD27 and its ligand CD70 is predominantly confined to lymphocytes, reflecting their specialized role in the adaptive immune response.

---

#### Product Data



Human peripheral blood lymphocytes was stained with APC Anti-Human CD27 clone O323 (color-filled histogram) or an isotype control (gray histogram).