

FITC Anti-human CD3 Antibody

Catalog Number:	109013, 109014
Size:	25 tests, 100 tests
Target Name:	CD3, T3, CD3ε
Regulatory Status:	RUO

PRODUCT DETAILS

Clone:	UCHT1
Application:	Flow Cytometry
Reactivity:	Human
Format:	FITC
Isotype:	Mouse IgG1
Antibody Type:	Monoclonal
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA
Protein Concentration:	Supplied at a lot-specific concentration.
Storage&Handling:	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Recommended Usage:	For flow cytometric staining, it is recommended to use 5 µL of this reagent per 0.5-1.0 million cells in a 100 µL volume. Optimal reagent performance should be determined by titration for each specific application. FITC has an excitation max at 493 nm and an emission max at 525 nm.
Excitation Laser:	Blue Laser (488 nm)
Isotype Control:	301415

BACKGROUND INFORMATION

CD3 is a critical component of the T-cell receptor (TCR) complex, essential for T-cell activation and the adaptive immune response. It is present on nearly all mature T lymphocytes and serves as a marker for their identification. Functionally, CD3 transduces activation signals from the TCR upon antigen recognition, triggering cascades that lead to cytokine production, cell proliferation, and differentiation. This process enables T-cells to respond effectively to pathogens and abnormal cells, anchoring immune defense mechanisms.


Structurally, CD3 is not a single protein but a multimeric complex composed of four distinct polypeptide chains: CD3γ, CD3δ, CD3ε, and CD3ζ. These chains associate non-covalently with the antigen-specific α and β chains of the TCR. Each CD3 subunit contains one or more immunoreceptor tyrosine-based activation motifs (ITAMs) in their cytoplasmic tails. Upon TCR engagement with an antigen-major histocompatibility complex (MHC), the ITAMs become phosphorylated, serving as docking sites for signaling molecules such as ZAP-70, which propagate downstream signaling required for full T-cell activation. CD3 itself does not bind directly to antigens; instead, it functions as a signaling adaptor within the TCR complex. Its effective operation relies on interactions

between the TCR and peptide-MHC complexes presented by antigen-presenting cells. The physical and functional association between CD3 and the TCR ensures that antigen recognition is tightly coupled to intracellular signaling pathways that determine T-cell fate and function.

Alterations in CD3 expression or signaling are implicated in various immune-related diseases. Mutations in CD3 genes can cause severe combined immunodeficiency (SCID), characterized by defective T-cell development. Dysregulated signaling may also contribute to autoimmune conditions, where inappropriate T-cell activation leads to tissue damage. Additionally, CD3 surface expression levels are sometimes altered in chronic infections and leukemias.


Therapeutically, CD3 is an important target in immunomodulation. Monoclonal antibodies such as muromonab-CD3 (OKT3) were among the first biologics used to prevent organ transplant rejection by depleting or inactivating T-cells. More recently, bispecific T-cell engagers (BiTEs) like blinatumomab exploit CD3 to redirect cytotoxic T-cells toward tumor cells expressing specific antigens, offering potent anti-cancer activity. By harnessing or modulating CD3-mediated signaling, modern immunotherapies continue to expand the therapeutic potential of T-cell-based treatments.

PRODUCT DATA



Human peripheral blood lymphocytes was stained with FITC Anti-Human CD3 clone UCHT1 (color-filled histogram) or an isotype control (gray histogram).

This product is supplied subject to the terms and conditions at www.innocyto.com/web/terms.php and may only be used as provided in the stated terms. Products are for Research Use Only.



Human peripheral blood lymphocytes stained with UCHT1 FITC and APC anti-human CD8 clone SK1

This product is supplied subject to the terms and conditions at www.innocyto.com/web/terms.php and may only be used as provided in the stated terms. Products are for Research Use Only.